



Which cardiovascular risks are associated
with insulin resistance in type 2 diabetes?

Site Map Your Account Support About Us Marketplace
Offerings: [Medscape.com](#) [Charts](#) [Mobile](#) [Logician](#) [CBSHealthwatch](#)

TECH CENTER 1600/2900

OCT 18 2001

RECEIVED

Medscape Medline Abstract



☐ Include in
selection list.

Document # 1 of 200 from MEDLINE

Relevance to Query:

Maximum Relevance:

Studies on the metabolism of [1-14C]5,8,11-eicosatrienoic (Mead) acid in rat hepatocytes.

Biochim Biophys Acta 1995 Oct 26;1259(1):82-8 (ISSN: 0006-3002)

Retterstol K; Woldseth B; Christophersen BO [[Find other articles with these Authors](#)]
Institute of Clinical Biochemistry, University of Oslo, Rikshospitalet, Norway.

The oxidation, esterification and formation of chain elongated and desaturated products of [1-14C] 5,8,11-eicosatrienoic (Mead) acid was studied. Liver cells from essentially fatty acid deficient (EFAD) and control rats were used. The metabolism of [1-14C]20:4, n-6 and [1-14C]20:5, n-3 were studied under the same experimental conditions. More 20:3, n-9 than 20:4, n-6 and 20:5, n-3 was oxidised both in EFAD and control cells. 20:3, n-9 was elongated to [14C]22:3, n-9 in both cell types and significant amounts of [14C]22:4, n-9 were formed in EFAD cells. Less 20:3, n-9 was esterified in phospholipids and more in triacylglycerol than observed with 20:4, n-6 and 20:5, n-3 in both cell types. 20:3, n-9 was mainly esterified in phosphatidylcholine and little was esterified in phosphatidylethanolamine compared to 20:4, n-6 and 20:5, n-3. In comparison, 20:3, n-9 was rather efficiently esterified in phosphatidylinositol as 18:0-20:3. [14C]22:4, n-9 formed from 20:3, n-9 in EFAD hepatocytes was esterified in triacylglycerol, not in phospholipids, unlike [14C]22:5, n-6 and [14C]22:6, n-3 which were mainly esterified in phospholipids.

Major Subject Heading(s)	Minor Subject Heading(s)	CAS Registry / EC Numbers
<ul style="list-style-type: none"> • 8,11,14-Eicosatrienoic Acid [analogs & derivatives] • Liver [metabolism] <p>Find other articles with similar Subjects.</p>	<ul style="list-style-type: none"> • 8,11,14-Eicosatrienoic Acid [metabolism] • Carbon Radioisotopes • Cells, Cultured • Fatty Acids, Essential [deficiency] • Kinetics • Oxidation-Reduction • Phospholipids [metabolism] • Radioisotope Dilution Technique 	<ul style="list-style-type: none"> • 0 (Carbon Radioisotopes) • 0 (Fatty Acids, Essential) • 0 (Phospholipids) • 0 (Triglycerides) • 2751-14-6 (mead acid) • 7324-41-6 (8,11,14-Eicosatrienoic Acid)